

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Principal Component Analysis

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- in-situ measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)




clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005




clear differential signal from VRS at low chl a, no signal over land


Monthly Average of Global Phytoplankton Conc. in July 2003 around the African Continent



VRS Fit Factor from SCIAMACHY from Vountas et al. 2005



VRS Fit Factor vs. Chl a from Vountas et al. 2005



With increasing chl a, VRS decreases rapidly high sensitivity of VRS-fit at low chl a

Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied

Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. *Polar Biology* 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Einbeziehung von Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: *Air Monitoring by Spectroscopic Techniques*, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. *Atmospheric Chemistry and Physics* 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to *Applied Optics* in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Principal Component Analysis

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- *in-situ* measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)




clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005




clear differential signal from VRS at low chl a, no signal over land


Monthly Average of Global Phytoplankton Conc. in July 2003 around the African Continent



VRS Fit Factor from SCIAMACHY from Vountas et al. 2005



VRS Fit Factor vs. Chl a from Vountas et al. 2005



With increasing chl a, VRS decreases rapidly high sensitivity of VRS-fit at low chl a

Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied

Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. Polar Biology 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Einbeziehung von Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: Air Monitoring by Spectroscopic Techniques, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. Atmospheric Chemistry and Physics 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to Applied Optics in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Principal Component Analysis

Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- in-situ measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)



clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005



clear differential signal from VRS at low chl a, no signal over land

Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied

Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. Polar Biology 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Einbeziehung von Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: Air Monitoring by Spectroscopic Techniques, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. Atmospheric Chemistry and Physics 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to Applied Optics in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Principal Component Analysis

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- in-situ measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)




clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005




clear differential signal from VRS at low chl a, no signal over land

Monthly Average of Global Phytoplankton Conc. in July 2003 around the African Continent




VRS Fit Factor from SCIAMACHY from Vountas et al. 2005



With increasing chl a, VRS decreases rapidly high sensitivity of VRS-fit at low chl a

VRS Fit Factor vs. Chl a from Vountas et al. 2005



Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied

Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. Polar Biology 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Einbeziehung von Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: Air Monitoring by Spectroscopic Techniques, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. Atmospheric Chemistry and Physics 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to Applied Optics in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Principal Component Analysis

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- *in-situ* measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)



clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005




clear differential signal from VRS at low chl a, no signal over land

Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied


Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Monthly Average of Global Phytoplankton Conc. in July 2003 around the African Continent



VRS Fit Factor from SCIAMACHY from Vountas et al. 2005




With increasing chl a, VRS decreases rapidly high sensitivity of VRS-fit at low chl a

Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates

Global Phytoplankton Concentrations averaged over July 2003



Good qualitative agreement between SCIAMACHY phytoplankton slant column and inverse VRS fit factor, and MERIS algal_1 chl a Below 35°S SZA at SCIAMACHY July data to high → here signal to noise ratio to bad

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. *Polar Biology* 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: *Air Monitoring by Spectroscopic Techniques*, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. *Atmospheric Chemistry and Physics* 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to *Applied Optics* in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Principal Component Analysis

Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- in-situ measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)



clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005



clear differential signal from VRS at low chl a, no signal over land

Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied

Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. Polar Biology 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Einbeziehung von Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: Air Monitoring by Spectroscopic Techniques, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. Atmospheric Chemistry and Physics 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to Applied Optics in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.

Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy on SCIAMACHY data and comparisons to MERIS



A. Bracher, M. Vountas, T. Dinter, B. Sierk, J. P. Burrows

University of Bremen, FB 1, Institute of Environmental Physics, P O Box 330440, D-28334 Bremen, Germany;
FAX: 0421-218-4555, email: bracher@uni-bremen.de



Global primary production & phytoplankton absorption & chl a

Objectives


Global models on marine primary production (PPR):

- function of fixed org. carbon to biomass (chl a) & light
- rarely consider spectral dependency of photosynthesis
- use ocean color satellite sensors (MERIS, MODIS, SeaWiFS,...) data on chl a, surface water reflectance and light penetration depth

Limitations to PPR models and chl a from ocean color:

- No global data set of *in situ* phytoplankton absorption spectra are available
- Slow sensitivity at low chl a conc.
- shapes of phytoplankton or pigment absorption are not constant (see figure to the right) due to different photosynthetic pigment composition & packaging

Phytoplankton Absorption in Different Biogeochemical Provinces of Atlantic Southern Ocean in Summer 95/96



Spectrum from Antarctic Circumpolar Current (ACC) used in SCIAMACHY DOAS retrieval explained below; Fig. from Bracher & Tilzer 2001

Specific absorption varies strongly for different provinces!

Important to derive information on attenuation of underwater components, above all the specific phytoplankton absorption from satellite data and Vibrational Raman Scattering (see Vountas et al. 2003) to improve chl a retrievals from satellite data and by that also global primary production estimates:

Synergistic use of the two ENVISAT instruments with information at 400 to 700 nm:

MERIS good spatial resolution ~1 km²

SCIAMACHY good spectral resolution ~0.2 - 0.4 nm

- Identify Vibrational Raman Scattering (VRS) and phytoplankton absorption in SCIAMACHY visible earth spectra by using the DOAS-method
- Determine phytoplankton slant columns and VRS fit factors and compare them to MERIS algal_1 chl a concentrations

DOAS-method used for ocean optics

Principal Component Analysis

Retrieval of trace gases (now applied to phytoplankton and VRS) by differential optical absorption spectroscopy (DOAS, Platt 1994) uses differential absorption signal of the molecular absorber in the earthshine spectrum. Forming the ratio of earthshine spectrum to extraterrestrial solar irradiance removes instrumental and Fraunhofer features.

Input:

Absorption cross section for molecular species: photosynthetic pigments*, yellow substance* (dissolved organic matter, DOM), O₃, NO₂, H₂O-vapor+ liquid* (measured)

Pseudo-absorber: spectrum of atmospheric Ring and VRS in liquid water* (modelled)

*contribute to the inherent optical properties of the water body, as seen in figure to the right


Procedure:

Least squares fit of DOAS equation on Beer's law to observation

Separation of high- and low frequency absorption features by low order polynomial

Output: Slant column density SCD = no. of molec. along average photon path or fit factor

Inherent Optical Properties for Case-1 Waters as Used in DOAS Retrievals of Phytoplankton Slant Columns and VRS



Problems of direct DOAS fitting of phytoplankton

- high correlation of polarisation features and ozone absorption in Chappius band of ozone
- phytoplankton column not zero over oligotrophic waters if solar background is used
- *in-situ* measured cross sections seem to be influenced by liquid water absorption

Solution: Principle Component Analysis PCA


(see Dinter 2005, Vountas et al. 2005)

- Eigenvector decomposition of large dataset of fit residuals
- first principle component included as pseudo-absorber and accounts for 99% of residual effects

SCIAMACHY chl a retrievals and Comparisons to MERIS


Conclusions

DOAS-Fit from 430 to 500 nm with PCA using Absorption Spectrum of *in situ* Phytoplankton (dominated by diatoms and <20µm from Bracher and Tilzer 2001, see first Figure)



clear differential signal from photosynthetic pigments

DOAS-Fit from 342 to 390 nm with PCA using Modelled Vibrational Raman Scattering (VRS) Reference Spectrum from Vountas et al. 2005




clear differential signal from VRS at low chl a, no signal over land

Phytoplankton absorption and VRS were identified in SCIAMACHY spectra when the DOAS method was applied


Both, SCIAMACHY global phytoplankton slant columns and VRS fit factor from DOAS-retrievals compare well to MERIS chl a

At low chl a SCIAMACHY VRS fit DOAS-retrievals are useful in open ocean waters (case-1) in addition to other Ocean Color data

Monthly Average of Global Phytoplankton Conc. in July 2003 around the African Continent




VRS Fit Factor from SCIAMACHY from Vountas et al. 2005



With increasing chl a, VRS decreases rapidly high sensitivity of VRS-fit at low chl a

VRS Fit Factor vs. Chl a from Vountas et al. 2005




Outlook

SCIAMACHY DOAS phytoplankton absorption and Vibrational Raman Scattering fits have the potential to

- improve the retrieval of marine chl a conc. and other oceanographic parameters from satellite data in case-1 waters at regional and local scales
- improve the retrieval of atmospheric trace gas concentrations from UV-VIS satellite spectra
- determine various specific phytoplankton absorption spectra which require validation with *in situ* measured spectra and distinguish different types of phytoplankton groups by spectral shape in conjunction with MERIS Ocean Color data
- provide parameters for global and regional marine primary production estimates


Global Phytoplankton Concentrations averaged over July 2003

MERIS algal_1 level-3 product




http://www.enviport.org/meris

SCIAMACHY Phytoplankton Slant Columns



SCIAMACHY Vibrational Raman Scattering Fit



Good qualitative agreement between SCIAMACHY phytoplankton slant column and inverse VRS fit factor, and MERIS algal_1 chl a Below 35°S SZA at SCIAMACHY July data to high → here signal to noise ratio to bad

References & Acknowledgements

Bracher A.U., Tilzer M.M. (2001) Underwater light field and phytoplankton absorbance in different surface water masses of the Atlantic Sector of the Southern Ocean. *Polar Biology* 24: 687-696.
Dinter T. (2005) Modellierung ozeanischer Rückstreuung unter Vibrations-Raman-Streuung und die Auswertung anhand von Satellitendaten. Diploma Thesis, University of Bremen, Bremen, Germany.
Platt U. (1994) Differential optical absorption spectroscopy (DOAS). In: *Air Monitoring by Spectroscopic Techniques*, edited by: Sigrist M.W., Chemical Analysis Series 127: 27-64.
Vountas M., Richter A., Witrock F., Burrows J.P. (2003) Inelastic scattering in ocean water and its impact on trace gas retrievals from satellite data. *Atmospheric Chemistry and Physics* 3: 1365-1375.

Vountas M., Dinter T., Sierk B., Bracher A., Burrows J.P. (2005) Chlorophyll retrieval studies using Differential Optical Absorption Spectroscopy. Submitted to *Applied Optics* in Oct. 2005

We thank ESA and DLR for providing us with SCIAMACHY level-1 data, and ESA and ACRI for MERIS algal_1 level-3 products. This work is funded by BMBF (SADOS 50/E0027), the University of Bremen and the State Bremen.